



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

A LIST OF THE BIRDS COLLECTED ON THE
ISLAND OF TOBAGO, WEST INDIES, BY W. W.
BROWN, JR., DURING APRIL
AND MAY, 1892.

BY CHARLES B. CORY.

THE collection made by Mr. Brown contained over seven hundred bird skins and a few mammals, including two specimens of *Sciurus griseogenys* Gray. The birds are especially interesting, several of them being new to the island.

- | | |
|--|--|
| <i>Sterna elegans</i> Gamb. | <i>Pachyrhamphus albo-griseus</i> Scf. |
| <i>Ardea virescens</i> Linn. | <i>Pachyrhamphus niger</i> Spix. |
| <i>Ionornis martinica</i> (Linn.). | <i>Chiroxiphia pareola</i> (Linn.). |
| <i>Ortalis ruficauda</i> Fard. | <i>Tyrannus melancholicus</i> Vieill. |
| <i>Columba rufina</i> (Temm.). | <i>Myiarchus tyrannulus</i> (Müll.). |
| <i>Engyptila verreauxi</i> (Bp.). | <i>Elainea pagana</i> (Licht.). |
| <i>Columbigallina rufipennis</i> (Bp.). | <i>Ostinops decumanus</i> (Pall.). |
| <i>Geotrygon linearis</i> (Knip et Prev.). | <i>Molothrus atronitens</i> (Cab.). |
| <i>Buteo latissimus</i> Wils. | <i>Euetheia bicolor</i> (Linn.). |
| <i>Spizaetus manduetyi</i> Daud. | <i>Spermophila lineata</i> (Gm.). |
| <i>Amazona amazonica</i> (Linn.). | <i>Volatinia splendens</i> (Vieill.). |
| <i>Crotophaga ani</i> Linn. | <i>Tanagra sclateri</i> Berlepsch. |
| <i>Dendrobates kirki</i> (Malh.). | <i>Tachyphonus melaleucus</i> Sparrm. |
| <i>Melanerpes terricolor</i> Berlepsch. | <i>Vireo chivi</i> (Vieill.). |
| <i>Chloronerpes rubiginosus</i> (Sw.). | <i>Hylophilus insularis</i> Sclater. |
| <i>Trogon collaris</i> Vieill. | <i>Progne dominicensis</i> (Gmel.). |
| <i>Momotus swainsoni</i> Sclater. | <i>Cæreba luteola</i> (Cab.). |
| <i>Ceryle americana</i> Gm. | <i>Arbelorhina cyanea</i> (Linn.). |
| <i>Galbula ruficauda</i> (Cuv.). | <i>Troglodytes tobagensis</i> Lawr. |
| <i>Chrysolampis moschitus</i> (Linn.). | <i>Thryothorus rutilus</i> (Vieill.). |
| <i>Florisuga mellivora</i> (Linn.). | <i>Mimus gilvus</i> Vieill.—In a series of |
| <i>Amazilia erythronota</i> (Less.). | <i>Mimus</i> from Tobago all the speci- |
| <i>Campylopterus ensipennis</i> (Swain.) | mens are paler and more ashy in |
| <i>Formicivora intermedia</i> Licht. | color on the upper parts than birds |
| <i>Dysithamnus semicinereus</i> Sclater. | from Grenada and St. Vincent. |
| <i>Thamnophilus doliatus</i> (Linn.). | <i>Merula xanthosceles</i> (Fard.). |
| <i>Dendroornis sussurans</i> Fard. | <i>Merula gymnophthalma</i> (Caban.). |